arXiv:1602.06901v3 [math.RA] 8 Jul 2016

Symbol Length of p-Algebras of Prime Exponent

Adam Chapman

Department of Computer Science, Tel-Hai College, Upper Galilee, 12208 Israel

Abstract

We prove that if the maximal dimension of an anisotropic homogeneous polynomial form of prime degree p over a field F with $\mathrm{char}(F) = p$ is a finite integer d greater than 1 then the symbol length of p-algebras of exponent p over F is bounded from above by $\left\lceil \frac{d-1}{p} \right\rceil - 1$, and show that every two tensor products of symbol algebras of lengths k and ℓ with $(k+\ell)p \geq d-1$ can be modified so that they share a common slot. For $p = 2$, we obtain an upper bound of $\frac{u(F)}{2} - 1$ for the symbol length, which is sharp when $I_q^3 F = 0$.

Keywords: Central Simple Algebras, p-Algebras, Symbol Length, Linkage, Homogeneous Polynomial Forms, Quadratic Forms, u-invariant
2010 MSC: 16K20 (primary); 11E76, 11E81 (secondary)

1. Introduction

A p-algebra is a central simple algebra of degree p^m over a field F with $\mathrm{char}(F) = p$ for some prime number p and positive integer m.

It was proven in [Alb61, Chapter 7, Theorem 30] that every p-algebra of exponent p is Brauer equivalent to a tensor product of algebras of the form

$$[\alpha,\beta)_{p,F} = F\langle x,y : x^p - x = \alpha, y^p = \beta, yx - xy = y\rangle$$

for some $\alpha \in F$ and $\beta \in F^\times$. We call such algebras "symbol algebras". An equivalent statement was proven in [MS82] when $\mathrm{char}(F) \neq p$ and F contains primitive pth roots of unity.

This means the group ${}_pBr(F)$ is generated by symbol algebras. The minimal number of symbol algebras required in order to express a given A in ${}_pBr(F)$ is

Email address: adam1chapman@yahoo.com (Adam Chapman)

called the symbol length of A. The symbol length of ${}_pBr(F)$ is the supremum of the symbol lengths of all $A \in {}_pBr(F)$. This number gives an indication of how complicated this group is, and due to the significance of this group, the symbol length has received special attention in papers such as [Flo13] and [Mat16].

In [Mat16, Theorem 3.4] it was proven that when $\mathrm{char}(F) \neq p$, F contains primitive pth roots of unity and the maximal dimension of an anisotropic homogeneous polynomial form of degree p over F is a finite integer d, the symbol length of A is at most $\left\lceil \frac{d+1}{p} \right\rceil - 1$. The case of p-algebras was also considered in that paper, but only over C_m-fields, which involve assumptions on the greatest dimensions of anisotropic homogeneous polynomial forms of any degree, not only p.

We prove that when $\mathrm{char}(F) = p$ and the maximal dimension of an anisotropic form of degree p over F is a finite integer d greater than 1, the symbol length of A is at most $\left\lceil \frac{d-1}{p} \right\rceil - 1$. We obtain this bound by first proving that every two tensor products of symbol algebras $\bigotimes_{i=1}^{k} C_i$ and $\bigotimes_{i=1}^{\ell} D_i$ with $(k+\ell)p \geq d-1$ can be modified so that they share a common slot.

In the last section, we focus on the case of $p = 2$. We explain how the same argument holds in this case if we replace d with the u-invariant, i.e. the maximal dimension of an anisotropic nonsingular quadratic form over F, and why the upper bound $\frac{u(F)}{2} - 1$ for the symbol length is sharp when $I_q^3 F = 0$.

2. Preliminaries

We denote by $M_p(F)$ the $p \times p$ matrix algebra over F. For two given central simple algebras A and B over F, we write $A = B$ if the algebras are isomorphic. We recall most of the facts we need on symbol algebras. For general background on these algebras see [Alb61, Chapter 7].

We start with some basic symbol changes. In the following remark and lemmas, let p be a prime integer and F be a field with $\mathrm{char}(F) = p$. We say that a homogeneous polynomial form φ over F is anisotropic if it has no nontrivial zeros.

Remark 2.1. (a) $M_p(F) = [\alpha, 1)_{p,F} = [0, \beta)_{p,F}$ for every $\alpha \in F$ and $\beta \in F^\times$.

(b) $[\alpha, \beta)_{p,F} \otimes [\alpha, \gamma)_{p,F} = M_p(F) \otimes [\alpha, \beta\gamma)_{p,F}$.

(c) $[\alpha, \gamma)_{p,F} \otimes [\beta, \gamma)_{p,F} = M_p(F) \otimes [\alpha + \beta, \gamma)_{p,F}$.

(d) If $[\alpha, \beta)_{p,F}$ contains zero divisors then it is $M_p(F)$.

(e) Given $K = F[x : x^p - x = \alpha]$, we denote by $\mathrm{N}_{K/F}$ the norm form $K \to F$. This form is homogeneous of degree p over F of dimension $[K : F] = p$.

Lemma 2.2. *Consider* $[\alpha,\beta)_{p,F} = F\langle x, y : x^p - x = \alpha, y^p = \beta, yx - xy = y\rangle$ *for some* $\alpha \in F$ *and* $\beta \in F^\times$. *Then*

(a) $[\alpha,\beta)_{p,F} = [\alpha, \mathrm{N}_{F[x]/F}(f)\beta)_{p,F}$ *for any* $f \in F[x]$ *with* $\mathrm{N}_{F[x]/F}(f) \neq 0$.

(b) $[\alpha,\beta)_{p,F} = [\alpha+\beta,\beta)_{p,F}$.

(c) $[\alpha,\beta)_{p,F} = [\alpha + v^p - v,\beta)_{p,F}$ *for any* $v \in F$.

(d) For any $v \in F$ *with* $\beta + v^p \neq 0$, $[\alpha,\beta)_{p,F} = [\alpha',\beta + v^p)_{p,F}$ *for some* $\alpha' \in F$.

Proof. Let f be an element in $F[x]$ with $\mathrm{N}_{F[x]/F}(f) \neq 0$. Then $(fy)^p = \mathrm{N}_{F[x]/F}(f)y^p = \mathrm{N}_{F[x]/F}(f)\beta$. Since $(fy)x - x(fy) = fy$, we have $[\alpha,\beta)_{p,F} = [x^p - x, (fy)^p)_{p,F} = [\alpha, \mathrm{N}_{F[x]/F}(f)\beta)_{p,F}$.

Write $z = x+y$. By [Cha15b, Lemma 3.1], z satisfies $z^p - z = x^p - x + y^p = \alpha + \beta$. Since $yz - zy = y$ we have $[\alpha,\beta)_{p,F} = [z^p - z, y^p)_{p,F} = [\alpha+\beta,\beta)_{p,F}$.

Write $t = x + v$. Then $t^p - t = x^p - x + v^p - v = \alpha + v^p - v$. Since $yt - ty = y$, we have $[\alpha,\beta)_{p,F} = [\alpha + v^p - v,\beta)_{p,F}$.

Let $v \in F$ be such that $\beta + v^p \neq 0$. Consider the element $y + v$. Since $(y+v)^p = \beta + v^p \in F^\times$, by [Alb61, Chapter 7, Lemma 10] we have $[\alpha,\beta)_{p,F} = [\alpha',\beta + v^p)_{p,F}$ for some $\alpha' \in F$. □

Note that by Lemma 2.2 (*a*), we have $[\alpha,\beta)_{p,F} = [\alpha,-\beta)_{p,F}$ for any symbol algebra $[\alpha,\beta)_{p,F}$.

Lemma 2.3. *Consider* $[\alpha,\beta)_{p,F}$ *and* $[\gamma,\delta)_{p,F}$ *for some* $\alpha,\gamma \in F$ *and* $\beta,\delta \in F^\times$. *Then*

(a) $[\alpha,\beta)_{p,F} \otimes [\gamma,\delta)_{p,F} = [\alpha+\gamma,\beta)_{p,F} \otimes [\gamma,\beta^{-1}\delta)$.

(b) If $\beta + \delta \neq 0$ *then* $[\alpha,\beta)_{p,F} \otimes [\gamma,\delta)_{p,F} = [\alpha+\gamma,\beta+\delta)_{p,F} \otimes C$ *for some symbol algebra* C *of degree* p.

Proof. Write $[\alpha,\beta)_{p,F} = F\langle x, y : x^p - x = \alpha, y^p = \beta, yx - xy = y\rangle$ and $[\gamma,\delta)_{p,F} = F\langle z, w : z^p - z = \gamma, w^p = \delta, wz - zw = w\rangle$. Note that the elements $x + z$ and y commute with the elements z and $y^{-1}w$, and that

$$(x+z)^p - (x+z) = x^p - x + z^p - z = \alpha + \gamma,$$

$$y(x+z)-(x+z)y = y,$$
$$(y^{-1}w)^p = y^{-p}w^p = \beta^{-1}\delta, \text{ and}$$
$$(y^{-1}w)z - z(y^{-1}w) = y^{-1}(wz-zw) = y^{-1}w.$$

Therefore $[\alpha,\beta)_{p,F} \otimes [\gamma,\delta)_{p,F} = F\langle x,y,z,w\rangle = F\langle x+z,y\rangle \otimes F\langle z,y^{-1}w\rangle = [\alpha+\gamma,\beta)_{p,F} \otimes [\gamma,\beta^{-1}\delta)_{p,F}$.

Now, the elements $x+z$ and $y+w$ satisfy

$$(y+w)^p = y^p + w^p = \beta+\delta, \text{ and}$$
$$(y+w)(x+z)-(x+z)(y+w) = y+w.$$

Therefore $F\langle x+z,y+w\rangle = [\alpha+\gamma,\beta+\delta)_{p,F}$. Hence $[\alpha,\beta)_{p,F} \otimes [\gamma,\delta)_{p,F} = [\alpha+\gamma,\beta+\delta)_{p,F} \otimes C$ for some central simple algebra C over F of degree p. Since C contains $y^{-1}w$ which satisfies $(y^{-1}w)^p \in F^\times$, C must be a symbol algebra ([Alb61, Chapter 7, Lemma 10]). □

3. Bounding the Symbol Length

Proposition 3.1. *Let p be a prime integer and let F be a field with* $\mathrm{char}(F) = p$. *Consider the algebras $[\alpha_1,\beta_1)_{p,F},\dots,[\alpha_k,\beta_k)_{p,F}$ for some $\alpha_1,\dots,\alpha_k \in F$ and $\beta_1,\dots,\beta_k \in F^\times$ where k is some positive integer. For each $i \in \{1,\dots,k\}$, write $[\alpha_i,\beta_i)_{p,F} = F\langle x_i,y_i : x_i^p - x_i = \alpha_i, y_i^p = \beta_i, y_ix_i - x_iy_i = y_i\rangle$.*

Let φ be the homogeneous polynomial form defined on

$$V = F \oplus F \oplus F[x_1] \oplus \cdots \oplus F[x_k]$$

by

$$\varphi(u,v,f_1,\dots,f_k) = u^p(\alpha_1+\cdots+\alpha_k) - u^{p-1}v + v^p + \mathrm{N}_{F[x_1]/F}(f_1)\beta_1 + \cdots + \mathrm{N}_{F[x_k]/F}(f_k)\beta_k.$$

(a) For every $(u,v,f_1,\dots,f_k) \in V$ with $u \neq 0$, we have $\bigotimes_{i=1}^k [\alpha_i,\beta_i)_{p,F} = \bigotimes_{i=1}^k C_i$ where $C_1,\dots,C_k$ are symbol algebras of degree p and $C_1 = [\varphi(1,\frac{v}{u},\frac{f_1}{u},\dots,\frac{f_k}{u}),\beta')_{p,F}$ for some $\beta' \in F^\times$.

(b) Let $(0,v,f_1,\dots,f_k) \in V$ such that the following elements are nonzero: $f_1,\dots,f_t$ for some $t \in \{1,\dots,k\}$, $\sum_{i=1}^s \mathrm{N}_{F[x_i]/F}(f_i)\beta_i$ for any $s \in \{1,\dots,t\}$ and $(\sum_{i=1}^t \mathrm{N}_{F[x_i]/F}(f_i)\beta_i) + v^p$. Then $\bigotimes_{i=1}^k [\alpha_i,\beta_i)_{p,F} = \bigotimes_{i=1}^k C_i$ where $C_1,\dots,C_k$ are symbol algebras of degree p and $C_1 = [\alpha', (\sum_{i=1}^t \mathrm{N}_{F[x_i]/F}(f_i)\beta_i) + v^p)_{p,F}$ for some $\alpha' \in F$.

(c) *If there exists* $(u, v, f_1, \ldots, f_k) \in V \setminus \{(0, \ldots, 0)\}$ *such that* $\varphi(u, v, f_1, \ldots, f_k) = 0$, *then* $\bigotimes_{i=1}^{k} [\alpha_i, \beta_i)_{p,F} = \bigotimes_{i=1}^{k} C_i$ *where* $C_1 = M_p(F)$ *and* $C_2, \ldots, C_k$ *are symbol algebras of degree* p.

Proof. Let $(u, v, f_1, \ldots, f_k) \in V$ with $u \neq 0$. For each $i \in \{1, \ldots, k\}$ with $\mathrm{N}_{F[x_i]/F}(\frac{f_i}{u}) \neq 0$, we apply Lemma 2.2 to change $[\alpha_i, \beta_i)_{p,F}$ to $[\alpha_i + \mathrm{N}_{F[x_i]/F}(\frac{f_i}{u})\beta_i, \mathrm{N}_{F[x_i]/F}(\frac{f_i}{u})\beta_i)_{p,F}$. Then for each $i \in \{2, \ldots, k\}$, we apply Lemma 2.3 (a) on the first and ith symbol algebras. The first algebra in the tensor product obtained after these modifications is $[\sum_{i=1}^{k}(\alpha_i + \mathrm{N}_{F[x_i]/F}(\frac{f_i}{u})\beta_i), \mathrm{N}_{F[x_1]/F}(\frac{f_1}{u})\beta_1)_{p,F}$ or $[\sum_{i=1}^{k}(\alpha_i + \mathrm{N}_{F[x_i]/F}(\frac{f_i}{u})\beta_i), \beta_1)_{p,F}$ depending on the value of $\mathrm{N}_{F[x_1]/F}(\frac{f_1}{u})$. By Lemma 2.2 (c) we can add $(\frac{v}{u})^p - \frac{v}{u}$ to the left slot of the first algebra. That proves part (a). If $\varphi(u, v, f_1, \ldots, f_k) = 0$ then also $\varphi(1, \frac{v}{u}, \frac{f_1}{u}, \ldots, \frac{f_k}{u}) = 0$, which means that the first algebra in the modified tensor product is a matrix algebra.

Let $(0, v, f_1, \ldots, f_k) \in V$ such that the following elements are nonzero: $f_1, \ldots, f_t$ for some $t \in \{1, \ldots, k\}$, $\sum_{i=1}^{s} \mathrm{N}_{F[x_i]/F}(f_i)\beta_i$ for any $s \in \{1, \ldots, t\}$ and $(\sum_{i=1}^{t} \mathrm{N}_{F[x_i]/F}(f_i)\beta_i) + v^p$. For each $i \in \{1, \ldots, t\}$ change $[\alpha_i, \beta_i)_{p,F}$ to $[\alpha_i, \mathrm{N}_{F[x_i]/F}(f_i)\beta_i)_{p,F}$. Then for each $i \in \{2, \ldots, t\}$ apply Lemma 2.3 (b) on the first and the ith symbol algebras. The first algebra in the resulting tensor product is $[\alpha_1 + \cdots + \alpha_t, \sum_{i=1}^{t} \mathrm{N}_{F[x_i]/F}(f_i)\beta_i)_{p,F}$. Then by Lemma 2.2 (d) we can change this algebra to $[\alpha', (\sum_{i=1}^{t} \mathrm{N}_{F[x_i]/F}(f_i)\beta_i) + v^p)_{p,F}$ for some $\alpha' \in F$. That proves part (b).

Let $(0, v, f_1, \ldots, f_k) \in V \setminus \{(0, \ldots, 0)\}$ such that $\varphi(0, v, f_1, \ldots, f_k) = 0$. If $f_1 = \cdots = f_k = 0$ then $\varphi(0, v, f_1, \ldots, f_k) = v^p = 0$ which is impossible for $v \in F^\times$. Therefore at least one f_i is nonzero. If there is one f_i which is nonzero and $\mathrm{N}_{F[x_i]/F}(f_i) = 0$ then the symbol algebra $[\alpha_i, \beta_i)_{p,F}$ contains a zero divisor and so it is a matrix algebra. Assume every nonzero f_i has $\mathrm{N}_{F[x_1]/F}(f_i) \neq 0$. Without loss of generality we can assume $f_1, \ldots, f_t$ are nonzero for some $t \in \{1, \ldots, k\}$ and $f_i = 0$ for $i > t$. Assume $\sum_{i=1}^{s} \mathrm{N}_{F[x_i]/F}(f_i)\beta_i = 0$ for some $s \in \{1, \ldots, t\}$. Assume s is minimal. Then we can change the tensor product so that the first algebra is $[\alpha', \sum_{i=1}^{s-1} \mathrm{N}_{F[x_i]/F}(f_i)\beta_i)_{p,F}$ and the sth algebra is $[\alpha_s, \mathrm{N}_{F[x_s]/F}(f_s)\beta_s)_{p,F}$. Since $\sum_{i=1}^{s-1} \mathrm{N}_{F[x_i]/F}(f_i)\beta_i = -\mathrm{N}_{F[x_s]/F}(f_s)\beta_s$, we have $[\alpha', \sum_{i=1}^{s-1} \mathrm{N}_{F[x_i]/F}(f_i)\beta_i)_{p,F} \otimes [\alpha_s, \mathrm{N}_{F[x_s]/F}(f_s)\beta_s)_{p,F} = [\alpha' + \alpha_s, \mathrm{N}_{F[x_s]/F}(f_s)\beta_s)_{p,F} \otimes M_p(F)$. Assume $\sum_{i=1}^{s} \mathrm{N}_{F[x_i]/F}(f_i)\beta_i \neq 0$ for all $s \in \{1, \ldots, t\}$. Then the tensor product can be changed such that the first algebra is $[\alpha', \sum_{i=1}^{t} \mathrm{N}_{F[x_i]/F}(f_i)\beta_i)_{p,F}$. This algebra contains a zero divisor because $\varphi(0, v, f_1, \ldots, f_k) = (\sum_{i=1}^{t} \mathrm{N}_{F[x_i]/F}(f_i)\beta_i) + v^p = 0$, which means it is a matrix algebra. That completes part (c). □

Theorem 3.2. *Let* p *be a prime integer and let* F *be a field with* $\mathrm{char}(F) = p$. *Assume the maximal dimension of an anisotropic homogeneous polynomial form*

of degree p over F is a finite integer d. Then every two tensor products $A = \bigotimes_{i=1}^{k}[\alpha_i, \beta_i)_{p,F}$ and $B = \bigotimes_{i=1}^{\ell}[\gamma_i, \delta_i)_{p,F}$ with $(k+\ell)p \geq d-1$ can be changed such that $\alpha_1 = \gamma_1$.

Proof. Let φ and ψ be the homogeneous polynomial forms of degree p as constructed in Proposition 3.1 for A and B respectively. If φ has a nontrivial zero then the first algebra in A can be assumed to be a matrix algebra, and so it can be written as $[\gamma_1, 1)_{p,F}$ and the statement follows. Similarly, the statement follows when ψ has a nontrivial zero.

Assume that both forms are anisotropic. The form $\varphi(u, v, f_1, \ldots, f_k) - \psi(u, 0, f_1', \ldots, f_\ell')$ is of dimension $(k+\ell)p+2$ over F. By assumption, $(k+\ell)p+2 \geq d+1$, which means that this form has a nontrivial zero, i.e. $\varphi(u, v, f_1, \ldots, f_k) = \psi(u, 0, f_1', \ldots, f_\ell')$ for some $u, v, f_1, \ldots, f_k, f_1', \ldots, f_\ell'$ such that not all of them are 0. If $u \neq 0$ then $\varphi(1, \frac{v}{u}, \frac{f_1}{u}, \ldots, \frac{f_k}{u}) = \psi(1, 0, \frac{f_1'}{u}, \ldots, \frac{f_\ell'}{u})$. By Proposition 3.1 (a) we can change both tensor products to have $\varphi(1, \frac{v}{u}, \frac{f_1}{u}, \ldots, \frac{f_k}{u})$ in the left slot of the first algebra.

Assume $u = 0$. At least one of the elements $v, f_1, \ldots, f_k, f_1', \ldots, f_\ell'$ is nonzero. If $f_i' = 0$ for every $i \in \{1, \ldots, \ell\}$ then $\psi(0, 0, f_1', \ldots, f_\ell') = 0$ and so $\varphi(0, v, f_1, \ldots, f_k) = 0$, but since φ is anisotropic we get $v = f_1 = \cdots = f_k = 0$, contradiction. Therefore $f_i' \neq 0$ for at least one i in $\{1, \ldots, \ell\}$. If $f_i = 0$ for every $i \in \{1, \ldots, k\}$ then $v^p = \psi(0, 0, f_1', \ldots, f_\ell')$, which means $0 = \psi(0, 0, f_1', \ldots, f_\ell') - v^p = \psi(0, -v, f_1', \ldots, f_\ell')$, i.e. ψ has a nontrivial zero, contradiction. Therefore $f_i \neq 0$ for at least one i in $\{1, \ldots, k\}$. By changing the order of the symbol algebras, we can assume $f_1, \ldots, f_t \neq 0$ for some $t \in \{1, \ldots, k\}$, $f_i = 0$ for every i with $i \geq t+1$, $f_1', \ldots, f_r' \neq 0$ for some $r \in \{1, \ldots, \ell\}$ and $f_i' = 0$ for every i with $i \geq r+1$. Both $(0, v, f_1, \ldots, f_k)$ and $(0, 0, f_1', \ldots, f_\ell')$ meet the requirements of Proposition 3.1 (b), and so we can change the first tensor product so that it has $\varphi(0, v, f_1, \ldots, f_k)$ in the second slot of the first symbol algebra and the second tensor product so that it has $\psi(0, 0, f_1', \ldots, f_\ell')$ in the second slot of the first symbol algebra. The statement then follows from [Cha15b, Theorem 3.2]. □

Corollary 3.3. *Let p be a prime integer and let F be a field with* $\mathrm{char}(F) = p$. *Assume the maximal dimension of an anisotropic homogeneous polynomial form of degree p over F is a finite integer d greater than 1. Then the symbol length of p-algebras of exponent p over F is bounded from above by $\left\lceil \frac{d-1}{p} \right\rceil - 1$.*

Proof. Write $n = \left\lceil \frac{d-1}{p} \right\rceil - 1$. Then it suffices to prove that for every tensor product $A = \bigotimes_{i=1}^{k}[\alpha_i, \beta_i)_{p,F}$ with $k \geq n+1$, $A = M_p(F) \otimes C_1 \otimes \cdots \otimes C_{k-1}$ for some symbol algebras $C_1, \ldots, C_{k-1}$ of degree p and the statement then follows by induction.

Note that if $k \geq n+1 = \left\lceil \frac{d-1}{p} \right\rceil$ then $kp \geq d-1$. Consider the algebra $[\alpha_1, \beta_1)_{p,F}$ and the tensor product $\bigotimes_{i=2}^{k} [\alpha_i, \beta_i)_{p,F}$. By Theorem 3.2, since $p(1+(k-1)) = kp \geq d-1$ we can assume $\alpha_1 = \alpha_2$, and so

$$[\alpha_1, \beta_1)_{p,F} \otimes [\alpha_2, \beta_2)_{p,F} \otimes \cdots \otimes [\alpha_k, \beta_k)_{p,F} = M_p(F) \otimes [\alpha_1, \beta_1\beta_2)_{p,F} \otimes [\alpha_3, \beta_3)_{p,F} \otimes \cdots \otimes [\alpha_k, \beta_k)_{p,F}.$$

□

Remark 3.4. By using similar methods, the upper bound for the symbol length of algebras of exponent p over a field F with $\mathrm{char}(F) \neq p$ containing primitive pth roots of unity appearing in [Mat16, Theorem 3.4] can be sharpened from $\left\lceil \frac{d+1}{p} \right\rceil - 1$ to $\left\lceil \frac{d-1}{p} \right\rceil - 1$ as well, where the maximal dimension of an anisotropic homogeneous polynomial form of degree p over F is d greater than 1. In the case of $p = 2$ and $\mathrm{char}(F) \neq 2$, d is the u-invariant of F, $u(F)$, and so this upper bound is $\left\lceil \frac{u(F)-1}{2} \right\rceil$ which appears in [Kah90, Theorem 2 (a)]. When we assume further that $I^3F = 0$, this bound is sharp (see [Kah90, Theorem 2 (b)]). For the construction of fields F with $\mathrm{char}(F) \neq 2$, $I^3F = 0$ and any even $u(F)$ see [Mer91, Theorem 4].

4. Quaternion Algebras

In this section we focus on the case of $p = 2$ and $\mathrm{char}(F) = 2$. In this case, the symbol algebras are quaternion algebras. The corresponding homogeneous polynomial forms of degree p constructed in the previous section are now quadratic forms. Quadratic forms in this case are of the shape

$$[a_1, b_1] \perp \cdots \perp [a_r, b_r] \perp \langle c_1, \ldots, c_t \rangle$$

for some $a_1, b_1, \ldots, a_r, b_r, c_1, \ldots, c_t \in F$. Each $[a_i, b_i]$ stands for the quadratic form $a_i u_i^2 + u_i v_i + b_i v_i^2$, $\langle c_1, \ldots, c_t \rangle$ is the diagonal form $c_1 w_1^2 + \cdots + c_t w_t^2$, and $\perp$ is the orthogonal sum. A quadratic form is nonsingular if $t = 0$. In particular, the dimension of a nonsingular quadratic form is always even. Note that $c[a, b]$ is isometric to $[\frac{a}{c}, bc]$ for $c \in F^\times$, so orthogonal sums of quadratic forms of the shape $c[a, b]$ are also nonsingular.

Given $K = F[x : x^2 + x = \alpha]$, the norm form $\mathrm{N}_{K/F} : K \to F$ mentioned in Remark 2.1 (e) is the quadratic form $[\alpha, 1]$.

The maximal dimension of an anisotropic quadratic form over F is denoted by $\hat{u}(F)$. The number d appearing in Theorem 3.2 and Corollary 3.3 is equal to $\hat{u}(F)$ when $p = 2$. The u-invariant of F, denoted by $u(F)$, is the maximal dimension of an anisotropic nonsingular quadratic form over F. Clearly $u(F) \leq \hat{u}(F)$, and

there are many examples in the literature where this inequality is strict (see for example [MTW91]). Therefore, using $u(F)$ instead of $\hat{u}(F)$ gives a better upper bound for the symbol length in the case of $p = 2$. We now rephrase Theorem 3.2 and Corollary 3.3 in terms of $u(F)$.

Theorem 4.1. *Let F be a field with* $\operatorname{char}(F) = 2$ *and $u(F) < \infty$. Then every two tensor products of quaternion algebras $A = \bigotimes_{i=1}^{k}[\alpha_i, \beta_i)_{2,F}$ and $B = \bigotimes_{i=1}^{\ell}[\gamma_i, \delta_i)_{2,F}$ with $(k + \ell)2 \geq u(F)$ can be changed such that $\alpha_1 = \gamma_1$.*

Proof. The proof is essentially the same as in Theorem 3.2. One just has to note that the form $\varphi(u, v, f_1, \ldots, f_k) - \psi(u, 0, f_1', \ldots, f_\ell')$ in this case is the quadratic form

$$[\alpha_1 + \cdots + \alpha_k + \gamma_1 + \cdots + \gamma_\ell, 1] \perp \beta_1[\alpha_1, 1] \perp \cdots \perp \beta_k[\alpha_k, 1] \perp \delta_1[\gamma_1, 1] \perp \cdots \perp \delta_\ell[\gamma_\ell, 1].$$

This quadratic form is nonsingular of dimension $(k+\ell)2+2$, which is greater than $u(F)$. Hence it has a nontrivial zero, and the proof continues in the same manner as the other proof. □

Corollary 4.2. *Let F be a field with* $\operatorname{char}(F) = 2$ *and $2 \leq u(F) < \infty$. Then the symbol length of algebras of exponent 2 over F is bounded from above by $\frac{u(F)}{2} - 1$.*

In certain cases this upper bound is sharp as Proposition 4.5 suggests. Before stating this proposition, we recall a few facts about quadratic forms: The unique (up to isometry) isotropic nonsingular quadratic form of dimension 2 over F is the hyperbolic plane $\mathbb{H} = [0, 1]$. Every nonsingular quadratic form φ over F can be written uniquely as $\varphi_{\text{an}} \perp \underbrace{\mathbb{H} \perp \cdots \perp \mathbb{H}}_{m \text{ times}}$ for some anisotropic form φ_{an} and some nonnegative integer m called the Witt index of φ. Two forms φ and ϕ are Witt equivalent if their underlying anisotropic subforms φ_{an} and ϕ_{an} are isometric. The group of Witt equivalence classes of nonsingular quadratic forms over F with $\perp$ as the group operation is denoted by I_qF. The "Arf invariant" or "discriminant" of a nonsingular form $\varphi = [a_1, b_1] \perp \cdots \perp [a_r, b_r]$ is the class of $\sum_{i=1}^{r} a_i b_i$ in the additive group $F/\wp(F)$ where $\wp(F) = \{\lambda^2 + \lambda : \lambda \in F\}$. The subgroup of I_qF of forms with trivial discriminant is I_q^2F. The Clifford invariant maps $2r$-dimensional nonsingular forms $\varphi = [a_1, b_1] \perp \cdots \perp [a_r, b_r]$ with trivial discriminant to tensor products of $r - 1$ quaternion algebras

$$E(\varphi) = [a_1b_1, b_rb_1)_{2,F} \otimes \cdots \otimes [a_{r-1}b_{r-1}, b_rb_{r-1})_{2,F}.$$

For every tensor product T of $r - 1$ quaternion algebras there exists a nonsingular form φ of dimension $2r$ with trivial discriminant such that $E(\varphi) = T$ (see [Cha15a,

Proof of Theorem 4.1]). The Clifford invariant defines a group epimorphism from I_q^2F to ${}_2Br(F)$ whose kernel is exactly I_q^3F.

Lemma 4.3. *If* $\mathrm{char}(F) = 2$, $I_q^3F = 0$ *and* $4 \leq u(F) < \infty$ *then there exists an anisotropic nonsingular quadratic form with trivial discriminant of dimension* $u(F)$.

Proof. Let ϕ be an anisotropic nonsingular form of dimension $u(F)$. Denote its discriminant by δ. If $\delta \in \wp(F)$, we are done, so assume $\delta \notin \wp(F)$. Write $\psi = \phi \perp [\delta, 1]$. Since ψ is of dimension $u(F) + 2$, it is isotropic. The Witt index of ψ can be either 1 or 2. If it is 2 then $\psi = \psi_{\mathrm{an}} \perp \mathbb{H} \perp \mathbb{H} = \psi_{\mathrm{an}} \perp [\delta, 1] \perp [\delta, 1]$, and so $\phi = \psi_{\mathrm{an}} \perp [\delta, 1]$. Note that the discriminant of ψ_{an} is trivial. Since $I_q^3F = 0$, ψ_{an} is universal (see [BC15, Proof of Theorem 4.1]), which contradicts the assumption that ϕ is anisotropic. Consequently the Witt index of ψ must be 1. Therefore ψ_{an} is an anisotropic nonsingular form of dimension $u(F)$ with trivial discriminant. □

Remark 4.4. By [Kah90, Theorem 2] (*b*), $u(F)$ is even when $\mathrm{char}(F) \neq 2$, $I^3F = 0$ and $u(F) > 1$. One can therefore prove in a similar way to Lemma 4.3 that if $\mathrm{char}(F) \neq 2$, $I^3F = 0$ and $4 \leq u(F) < \infty$ then there exists an anisotropic quadratic form with trivial discriminant of dimension $u(F)$.

Proposition 4.5. *When* $\mathrm{char}(F) = 2$, $I_q^3F = 0$ *and* $2 \leq u(F) < \infty$, *the bound* $\frac{u(F)}{2} - 1$ *for the symbol length of algebras of exponent 2 over* F *is sharp.*

Proof. Write $u(F) = 2n$ for some positive integer n. The statement is clearly true when $n = 1$, so assume $n \geq 2$. It is enough to find an algebra of exponent 2 over F whose symbol length is $\frac{u(F)}{2} - 1 = n - 1$. By Lemma 4.3 there exists an anisotropic nonsingular form φ with trivial discriminant of dimension $2n$. We claim that the symbol length of $E(\varphi)$ is $n - 1$. Clearly it is no greater than $n - 1$. Suppose $E(\varphi)$ is Brauer equivalent to a tensor product T of k quaternion algebras where $k < n - 1$. Then there exists some nonsingular form ϕ of dimension $2(k+1)$ with trivial discriminant such that $E(\phi) = T$. Since $E(\phi)$ and $E(\varphi)$ are Brauer equivalent, the forms ϕ and φ are equivalent modulo I_q^3F. However, $I_q^3F = 0$, which means that ϕ and φ are Witt equivalent, but this is impossible because φ is anisotropic of dimension $2n$ and ϕ is of dimension $2(k + 1) < 2n$. □

Corollary 4.2 and Proposition 4.5 provide characteristic 2 analogues to parts (*a*) and (*b*) of [Kah90, Theorem 2]. For the construction of fields F with $\mathrm{char}(F) = 2$, $I_q^3F = 0$ and any even u-invariant see [EKM08, Theorem 38.4].

Remark 4.6. There is a typographical error in the first sentence, second paragraph of the proof of [BC15, Theorem 4.1]. The correct sentence should be: "Now we show that if $f \in I_q^2 F$ is an anisotropic form of dimension at least 6 and f_K is isotropic for some $K = F[x : x^2 + x = a]$ then $f_K = \mathbb{H} \perp f_0$ where $\mathbb{H}$ is a hyperbolic plane and f_0 is anisotropic".

Acknowledgments

The author thanks the anonymous referee for the helpful comments on the manuscript.

Bibliography

References

[Alb61] A. Adrian Albert, *Structure of algebras*, Revised printing. American Mathematical Society Colloquium Publications, Vol. XXIV, American Mathematical Society, Providence, R.I., 1961. MR 0123587 (23 #A912)

[BC15] Demba Barry and Adam Chapman, *Square-central and Artin-Schreier elements in division algebras*, Arch. Math. (Basel) **104** (2015), no. 6, 513–521. MR 3350341

[Cha15a] Adam Chapman, *Chain equivalences for symplectic bases, quadratic forms and tensor products of quaternion algebras*, J. Algebra Appl. **14** (2015), no. 3, 1550030, 9. MR 3275567

[Cha15b] ———, *Common subfields of p-algebras of prime degree*, Bull. Belg. Math. Soc. Simon Stevin **22** (2015), no. 4, 683–686. MR 3429179

[EKM08] Richard Elman, Nikita Karpenko, and Alexander Merkurjev, *The algebraic and geometric theory of quadratic forms*, American Mathematical Society Colloquium Publications, vol. 56, American Mathematical Society, Providence, RI, 2008. MR 2427530

[Flo13] Mathieu Florence, *On the symbol length of p-algebras*, Compos. Math. **149** (2013), no. 8, 1353–1363. MR 3103068

[Kah90] B. Kahn, *Quelques remarques sur le u-invariant*, Sém. Théor. Nombres Bordeaux (2) **2** (1990), no. 1, 155–161. MR 1061764

[Mat16] Eliyahu Matzri, *Symbol length in the Brauer group of a field*, Trans. Amer. Math. Soc. **368** (2016), no. 1, 413–427. MR 3413868

[Mer91] A. S. Merkur′ev, *Simple algebras and quadratic forms*, Izv. Akad. Nauk SSSR Ser. Mat. **55** (1991), no. 1, 218–224. MR 1130036 (93b:16025)

[MS82] A. S. Merkur′ev and A. A. Suslin, *K-cohomology of Severi-Brauer varieties and the norm residue homomorphism*, Izv. Akad. Nauk SSSR Ser. Mat. **46** (1982), no. 5, 1011–1046, 1135–1136. MR 675529 (84i:12007)

[MTW91] P. Mammone, J.-P. Tignol, and A. Wadsworth, *Fields of characteristic* 2 *with prescribed u-invariants*, Math. Ann. **290** (1991), no. 1, 109–128. MR 1107665